

**Metroplex Colon & Rectal Specialist**  
**Raymond Staniunas, M.D., FACS**  
**Board Certified Colon & Rectal Surgery**

Office Toll Free 1-855-273-2825  
Fax 817-701-0575

**DEFECOGRAPHY**

**PATIENT INFORMATION**

**WHAT IS DEFECOGRAPHY?**

Defecography, or evacuation proctography, is an X-ray test that shows the rectum and anal canal as they change during defecation (having a bowel movement). This test is used to evaluate for disorders of the lower bowel that are not evident by tests such as colonoscopy or sigmoidoscopy.

**PREPARATION FOR THE PROCEDURE**

- Give yourself a Fleet enema 2 hours prior to your study. Repeat enema after 15 minutes. You can purchase the two Fleet enemas from a pharmacy or supermarket.
- You should not eat anything during the two hours prior to the procedure. If you are diabetic, this may involve adjusting your diabetic medications.
- You should take your regular medications with a small sip of water at least 2 hours prior to the study.

**THE DEFECOGRAPHY TEST**

Defecography is a technique in which a barium contrast medium is introduced into your rectum after the radiologist performs a rectal examination. The barium is visible within the rectum on X-rays. During the test, you are instructed to defecate (empty the rectum) on a commode while X-rays of the pelvis are taken. These X-rays are taken while the person is sitting at rest, straining, squeezing, and during defecation. This type of test, although awkward, provides valuable information that may aid your doctor in diagnosing your problem.

**WHEN IS FECOGRAPHY USED?**

Defecation (having a bowel movement) is a complex action requiring coordination with relaxation and contraction of large number of muscles. It is controlled by the nervous system, but is also under voluntary control. The process of defecation is initiated by the arrival of stool into the rectum. This sensation leads to a chain of events which ends in evacuation of stool from the anus. The act of defecation is voluntarily controlled in healthy, normally functioning people.

The following is a list of some conditions for which defecography can be used to gather more information about a patient's condition and/or confirm a diagnosis.

- Chronic Constipation - evaluating for functional obstruction
- Rectal Prolapse
- Rectocele (an outpouching of the rectum)
- Fecal Incontinence
- Anismus (inappropriate spasm of the anal sphincter)

If you have any questions, please call our office at 1-855-273-2825.